

STRUCTURE of ATOM

1. The angular momentum of electron of H-atom is proportional to

- (a) r^2 (b) $1/r$ (c) $r\sqrt{}$ (d) $\frac{1}{\sqrt{r}}$

Answer: (c)

From Bohr's Model,

Centripetal force of electron = Columbic force of attraction on electron by nucleus

$$\frac{mv^2}{r} \propto \frac{e^2}{r^2}$$

$$\frac{mv^2}{r} \propto \frac{e^2}{r^2} \Rightarrow (mv^2)r \propto \frac{e^2}{r}$$

$$(mvr)^2 \propto e^2 mr$$

$$(\text{Angular momentum})^2 \propto r$$

Angular momentum $\propto \sqrt{r}$

2. The total energy in 1st orbit of hydrogen atom is given by

- (a) $\frac{2\pi^2 k^2 e^4 m}{h^3}$ (b) $-\frac{2\pi^2 k^2 e^2 m}{h^2}$
 (c) $\frac{2\pi^2 k^2 e^4 m}{h^2}$ (D) $\frac{2\pi^2 k^2 e^4 m}{h^2}$

Answer: (d)

• Energy of electron in Bohrs orbit $= -\frac{2\pi^2 k^2 z^2 e^4 m}{n^2 h^2}$

- for 1st orbit of hydrogen $n=1$
 $z=1$
- \therefore energy $= -\frac{2\pi^2 k^2 e^4 m}{h^2}$

3. If the radius of second Bohrs orbit of hydrogen atom is a_0 than the radius of third Bohrs orbit of Be^{3+} ion will be

- (a) $9a_0$ (b) $\frac{9a_0}{8}$
 (c) $\frac{9a_0}{16}$ (d) $\frac{9a_0}{4}$

Answer: (c)

- $r_H = 0.529 \frac{n^2}{Z^H} \text{ \AA}$ $r_{Be^{3+}} = 0.529 \frac{n^2}{Z^{Be^{3+}}}$
- $\frac{r_H}{r_{Be^{3+}}} = \frac{n^2}{Z^H} \cdot \frac{Z^{Be^{3+}}}{n^2} = \frac{(2)^2}{1} \cdot \frac{1}{4} = 1$
- $\frac{a_0}{r_{Be^{3+}}} = \frac{16}{9} \Rightarrow r_{Be^{3+}} = \frac{9a_0}{16}$

4. If the total energy of an electron in the 1st shell of H-atom is -13.6 eV then its potential energy in the 1st excited state would be:

- (a) $+6.8 \text{ eV}$ (b) $+20.4 \text{ eV}$ (c) -6.8 eV (d) $+3.4 \text{ eV}$

Answer: (c)

$$E_n = -\frac{13.6 Z^2}{n^2}$$

For Hydrogen atom, $Z = 1$

$$E_n = -\frac{13.6}{n^2}$$

$$E_1 = -13.6 \text{ eV}$$

$$E_2 = -3.4 \text{ eV}$$

Potential energy of an electron = 2 × Total energy of electron

$$(P.E.)_2 = 2E_2 = -6.8 \text{ eV}$$

5. If the speed of the electron in the first Bohr orbit of hydrogen atom be 'x' then the speed of electron in the third Bohr orbit is

(a) $\frac{x}{9}$

(b) $\frac{x}{3}$

(c) x

(d) 9x

Answer: (b)

The speed of electron in Bohr's orbit for hydrogen atom is given by

$$v = \frac{2\pi e^2 Z}{nh} = \frac{2.188 \times 10^8 Z}{n} \text{ cms}^{-1}$$

For the first Bohr orbit of H-atom,

$$Z = 1, n = 1$$

$$v_1 = 2.188 \times 10^8 \text{ cms}^{-1} = x$$

$$v_3 = \frac{2.188 \times 10^8}{3} \text{ cms}^{-1} = \frac{x}{3}$$

$$\text{Or } Z^2 = \frac{\Delta E}{\Delta E_n} = \frac{5.0 \times 10^{-18}}{1.63 \times 10^{-18}} = 3.06$$

$$z = 2$$

8. Possible number of nodal planes present in the Mth shell of H-like species are:

- (a)13 (b)14 (c)11 (d)9

Answer: (c)

Mth shell is n = 3

	3s	3px	3py	3pz	3dxy	3dyz	3dzx	3d _{xy²-y²}	3d _{z²}
Nodal Planes	0	1	1	1	2	2	2	2	0

Hence, Total no. of nodal planes

$$= 0 + 1 + 1 + 1 + 2 + 2 + 2 + 2 + 0 = 11$$

9. The no. of radial nodes for 3Px orbital is

- (a)One (b)Two (c)Three (d)Infinite

Answer: (a)

$$\text{No. of radial nodes} = n - l - 1 = 3 - 1 - 1 = 1$$

10. Photoelectric emission is observed from a surface for frequencies ν_1 and ν_2 of incident radiations ($\nu_1 > \nu_2$). If the maximum kinetic energy of photoelectrons in the two cases are in the ratio of 2 : 1, then threshold frequency ν_0 is given by

- (a) $\frac{\nu_2 - \nu_1}{2 - 1}$ (b) $\frac{2\nu_1 - \nu_2}{2 - 1}$ (c) $\frac{2\nu_2 - \nu_1}{2 - 1}$ (d) $\frac{\nu_2 - \nu_1}{2}$

Answer: (c)

$$h\nu = h\nu_0 + \text{K.E.}$$

$$\text{KE}_1 = h\nu_1 - h\nu_0$$

$$\text{KE}_2 = h\nu_2 - h\nu_0$$

$$\frac{\text{KE}_1}{\text{KE}_2} = \frac{h(\nu_1 - \nu_0)}{h(\nu_2 - \nu_0)} = \frac{2\nu_1 - \nu_0}{\nu_2 - \nu_0}; \nu_0 = \frac{2\nu_2 - \nu_1}{2 - 1}$$

11. The spin angular momentum for the 's' electron in H-atom

is

- (a) $\frac{\sqrt{3}h}{4\pi}$ (b) $\frac{h}{2\pi}$ (c) $\frac{h}{4\pi}$ (d) $\frac{\sqrt{3}h}{2\pi}$

Answer: (a)

Electronic configuration of Hydrogen atom is $1s^1$

Total spin of an unpaired electron, $S = \frac{1}{2}$

$$\therefore \text{Spin angular momentum} = S(S+1)\frac{h}{2\pi} = \sqrt{\frac{1}{2}(\frac{1}{2}+1)}\frac{h}{2\pi} = \frac{\sqrt{3}h}{4\pi}$$

12. If a dye absorbs a photon of wavelength λ and re-emits the absorbed energy into two photons of wavelengths λ_1 and λ_2 respectively. Then

(a) $\lambda = \frac{\lambda_1 + \lambda_2}{\lambda_1 \lambda_2}$ (b) $\lambda = \frac{\lambda_1 \lambda_2}{\lambda_1 + \lambda_2}$ (c) $\lambda = \frac{\lambda_2 \lambda_2}{\lambda_1 + \lambda_2}$ (d) $\lambda = \frac{\lambda_1 \lambda_2}{(\lambda_1 + \lambda_2)^2}$

Answer: (b)

$$E = E_1 + E_2$$

$$\frac{hc}{\lambda} = \frac{hc}{\lambda_1} + \frac{hc}{\lambda_2}$$

$$\frac{hc}{\lambda} = hc \left[\frac{\lambda_1 + \lambda_2}{\lambda_1 \lambda_2} \right]$$

$$\lambda = \frac{\lambda_1 \lambda_2}{\lambda_1 + \lambda_2}$$

13. An electron is allowed to move freely in a closed cubic box of length 10 cm. The minimum uncertainty in its velocity will be observed as

(a) $4 \times 10^{-3} \text{ m/s}$ (b) $5 \times 10^{-4} \text{ m/s}$ (c) $4 \times 10^{-5} \text{ m/s}$ (d) $4 \times 10^{-6} \text{ m/s}$

Answer: (b)

$$\Delta x = a\sqrt{3}, \Delta x = 10\sqrt{3}, \Delta x \cdot \Delta p = \frac{h}{4\pi}$$

$$\Delta V = \frac{h}{4\pi m \Delta x} = \frac{6.62 \times 10^{-34}}{4 \times 3.14 \times 9.1 \times 10^{-31} \times 10\sqrt{3}} \approx 5 \times 10^{-4} \text{ m/s}$$

14. If a_0 be the radius of first Bohr's orbit of H-atom, the de-Broglie's wavelength of an electron revolving in the third Bohr's orbit will be

- (a) $6\pi a_0$ (b) $4\pi a_0$
 (c) $2\pi a_0$ (d) πa_0

Answer: (a)(i)

$$\frac{mvrnh}{2\pi}$$

deBroglie equation = $Ph = \frac{h}{\lambda} = mv \dots(ii)$

placing the value of mv from (ii) and (i) for 3rd orbit

$$\frac{h}{\lambda} \cdot 3 = \frac{3h}{2\pi}$$

$$\Rightarrow \lambda = \frac{2\pi}{3}$$

$$\lambda = \frac{2\pi r_3}{3}$$

or $r = n^2 a_0 = 9a_0$

$$\text{so } \lambda = \frac{2\pi \cdot 9a_0}{3} = 6\pi a_0$$

15. The probability of finding an electron residing in a p_x orbital is zero

- (a) in yz plane (b) in xy plane
 (c) in the y direction (d) in the z direction

Answer: (a)

Nodal plane of P_x orbital is yz -plane.

16. In a single hydrogen atom, the electron is excited to its 6th orbit. The maximum no. of distinct lines possible, when it comes to the ground state is:

- (a)12 (b)15 (c)5 (d)22

Answer: (c)

Maximum number of spectral lines (or distinct lines) if an electron jumps from n_2 to n_1 monoelectronic species = $n_2 - n_1$.

If an electron jumps from $n_2=6$ to $n_1=1$ in a Hydrogen atom, then maximum number of spectral lines = $6 - 1 = 5$

17. The difference between n th and $(n+1)$ th Bohr's radius of H-atom is equal to (n) th Bohr's radius. Hence the value of 'n' is:

- (a)5 (b)2 (c)6 (d)4

Answer: (c)

Bohr's radius of n^{th} orbit, $r_n = \frac{0.529 n^2}{Z} \text{ \AA}$

$$r_{n+1} - r_n = r_{n-1}$$

$$\frac{0.529(n+1)^2}{Z} - \frac{0.529(n)^2}{Z} = \frac{0.529(n-1)^2}{Z}$$

$$0.529[(n+1)^2 - (n)^2] = 0.529(n-1)^2$$

$$(n+1)^2 - n^2 = (n-1)^2$$

$$n^2 + 1 + 2n - n^2 = n^2 + 1 - 2n$$

On solving, $n = 4$

18. Consider one He⁺ ion in excited state ($n = 5$). Which of the following observations will hold true as per the Bohr's model?

- (a) 10 emission spectral lines will be seen
 - (b) The ionization energy needed is less than 2 eV
 - (c) The longest emitted wavelength is less than $10/R$ ($R =$ Rydberg's constant)
 - (d) The electronic separation from the centre of nucleus is more than 60\AA
- Answer: (d)

(a) A multitude of He⁺ will produce 10 possible emission spectral lines. However, a single He⁺ can produce maximum 4 lines (not more)

$$5 \rightarrow 4 \rightarrow 3 \rightarrow 2 \rightarrow 1$$

(b) $E_n = -\frac{13.6}{n^2} = -13.6 \times \frac{1}{5^2} = -13.6 \times \frac{1}{25} = -2.176 \text{ eV}.$

$$\therefore 1E = + 2.176 \text{ eV} > 2 \text{ eV}$$

$$\frac{1}{\lambda} = RZ^2 \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$


$$(c) \frac{1}{\lambda} = R \cdot 4 \left[\frac{1}{4} - \frac{1}{9} \right]$$

$$\frac{1}{\lambda} = 4R \left[\frac{9}{16 \times 25} \right]$$

$$\lambda = \frac{16 \times 25}{4 \times 9 \times R} = \frac{100}{9R} > \frac{10}{R}$$

$$(d) r_n = 0.5 \frac{3n^2}{Z} = 0.53 \times \frac{5^2}{Z} = 6.625 \text{ \AA} > 6 \text{ \AA}$$

19. The given diagram indicates the energy levels of certain atom. When an electron moves from 2E level to E level, a photon of wavelength λ is emitted. The wavelength of photon emitted during its transition from $\frac{4E}{3}$ level to E level is:



$$(a) \frac{\lambda}{3}$$

$$(b) \frac{3\lambda}{4}$$

$$(c) \frac{4\lambda}{3}$$

$$(d) 3\lambda$$

Answer: (d)

$$h\nu = 2E - E$$

$$\frac{hc}{\lambda} = E \text{ or } \lambda = \frac{hc}{E}$$

When the electron jumps from $4E_3$ level to E level, a photon of wavelength λ_1 is emitted. Thereby,

$$\frac{hc}{\lambda_1} = \frac{4E}{3} - E = \frac{E}{3}$$

$$\lambda_1 = 3\lambda$$

20. Total number of electrons in Cu^{2+} for which the summation of azimuthal quantum number and magnetic quantum number is zero is

- (a) 4 or 8 (b) 11 or 12 (c) 5 or 6 (d) 10 or 11

Answer: (b)

Electronic configuration of Cu^{2+} is $1s^2 2s^2 2p^6 3s^2 3p^6 3d^9$

Electrons

$1s^2$	$\ell + m = 0$	2 [For s-subshell, $\ell = 0$ and $m = 0$]
$2s^2$	$\ell + m = 0$	2
$2p_x^2 2p_y^2 2p_z^2$	$\ell + m = 0$	2 [For p-subshell, $\ell = 1$ and $m = -1$]
$3s^2$	$\ell + m = 0$	2
$3p_x^2 3p_y^2 3p_z^2$	$\ell + m = 0$	2
$3d^9$	$\ell + m = 0$	1 or 2 [For d-subshell, $\ell = 2$ and $m = -2$]

21. Photons having energy equivalent to binding energy of 4th state of He^+ ion is used on the metal surface of work function 1.4 eV. If electrons are further accelerated through

the potential difference of 4V then the minimum value of de-Broglie wavelength associated with the electron is

- (a) 1.1 Å (b) 9.15 Å (c) 5 Å (d) 11 Å

Answer: (c)

$$\text{Binding energy of He}^+ = 13.6 \times \frac{2^2}{4} = \frac{13.6}{4} = 3.4 \text{ eV}$$

$$\text{Total K.E. of electron} = 3.4 - 1.4 + 4 \text{ eV} = 6 \text{ eV}$$

$$\lambda = \frac{h}{\sqrt{2meV}} = \frac{h}{\sqrt{2m \times 6}} = 5 \text{ Å}$$

22. If Rutherford would have used β -particles instead of α -particles in his Gold-leaf experiment, which of the following observations would definitely not have been made:

- (a) β -particles scattering beyond the gold foil.
- (b) Majority of β -particles passing undeflected through the foil.
- (c) Very few β -particles getting absorbed.
- (d) Wave nature of electron

Answer: (a)

α -particles are scattered by positively charged nucleus; β -particles will converge while passing through atom as they bear negative charge.

23. A 1-kW radio transmitter operates at a frequency of 880 Hz. How many quantas per second does it emit?

- (a) 1.71×10^{21} (b) 1.71×10^{30} (c) 6.02×10^{23} (d) 2.85×10^{26}

Answer: (b)

A 1kW radio transmitter emits 1000 Joules of radio waves per second.

Energy of one quanta, $E = h\nu$

$$= 6.626 \times 10^{-34} \times 880 \times 10^3$$

$$= 5.83 \times 10^{-28} \text{J}$$

Number of quantas emitted per second = $\frac{\text{Energy emitted per sec}}{\text{Energy of one quanta}}$

$$= \frac{1000}{5.83 \times 10^{-28}} = 1.71 \times 10^{30}$$

24. The ratio of the e/m values of a proton and an α -particle is

- (a) 2 : 1 (b) 1 : 1 (c) 1 : 2 (d) 1 : 4

Answer: (a)

For a proton $(H)_e = \frac{1.6 \times 10^{-19} \text{C}}{1.67 \times 10^{-27} \text{kg}}$

For an α -particle $(He\ 2)_e = \frac{2 \times 1.6 \times 10^{-19} \text{C}}{4 \times 1.67 \times 10^{-27} \text{kg}}$

$$\frac{(e/m)_p}{(e/m)_\alpha} = \frac{4}{2} = \frac{2}{1}$$

25. For sets of values of quantum numbers (n, l, m and s) are given below. Which of these does not provide a permissible solution of the wave equation?

- (a) $3, 2, -\frac{2}{2}, 1$ (b) $3, 3, 1, \frac{1}{2}, -1$ (c) $3, 2, \frac{1}{2}, 1$ (d) $3, 1, 1, \frac{1}{2}, -1$

Answer: (b)

$n = l$ is not permissible.

Value of l is always less than the value of n .

26. Angular momentum of an electron in a Bohr's orbit of He^+ is $3.1652 \times 10^{-34} \text{ kg.m}^2/\text{sec}$ The wave number of the spectral line emitted when an electron falls from this level to the first excited state is.

- (a) $3R$ (c) $3R$ (b) $\frac{5R}{9}$
 $\frac{4}{4}$ (d) $\frac{8R}{9}$

Answer: (b)

$$3.1652 = \frac{nh}{2\pi}$$

$$n \times 6.626 \times 10^{-34}$$

$$3.1652 = \frac{\quad}{2 \times \frac{22}{7}}$$

$n = 3$


$$v = R \cdot Z^2 \left(\frac{1}{2^2} - \frac{1}{1^2} \right)$$

$$v = R \cdot 2^2 \left(\frac{1}{2^2} - \frac{1}{1^2} \right) = \frac{5R}{9}$$

27. The electronic configuration of Fe³⁺ is

- (a) [Ar]3d⁵4s¹ (b) [Ar]3d⁶4s⁰ (c) [Ar]3d⁵4s⁰ (d) [Ar]3d⁶4s²

Answer: (c)



28. The value of the magnetic moment of a particular ion is 2.83 Bohr magneton. The ion is

- (a) Fe²⁺ (b) Ni²⁺ (c) Mn²⁺ (d) Co³⁺

Answer: (b)

The spin only magnetic moment is given by ,

$$\mu_s = \sqrt{n(n+2)} \text{ BM}$$

Where 'n' is no. of unpaired electrons.

$$2.83 = \sqrt{n(n+2)} \Rightarrow n=2$$

The value of $\mu = 2.83 \text{ BM}$ corresponds to the presence of two unpaired electrons. So the ion is Ni²⁺(3d⁸).

29. The ratio of the radii of the first three Bohr orbits is :

- (a) 1:05:0.33 (b) 1:2:3 (c) 1:4:9 (d) 1:8:27

Answer: (c)

$$r_n = r_1 \times n^2$$

For first orbit $r_1 = r_1$

For second orbit, $r_2 = r_1 \times 2^2 = 4r_1$

For third orbit, $r_3 = r_1 \times 3^2 = 9r_1$

Hence the ratio of radii of Bohr's orbits is 1:4:9.

30. Atomic number of chromium is 24, then Cr^{3+} will be:

- (a) diamagnetic (b) paramagnetic
(c) ferromagnetic (d) none of these

Answer: (b)

Electronic configuration of Cr^{3+} is $1s^2 2s^2 2p^6 3s^2 3p^6 3d^3$

Cr^{3+} has three unpaired 3d electrons. Hence, Cr^{3+} will be paramagnetic.

31. Find the incorrect set of quantum number.

(a) $n=2, l=1, m=0, s=1/2$ (b) $n=3, m=3, s=+1/2$

(c) $n=5, l=3, m=2, s=+1/2$ (d) $n=4, l=2, m=3, s=+1/2$

(d)

Answer: (d) $n=3, l=$

If the angular momentum quantum number of an electron $l=3$, which means $(n-1)$ values are possible for l , so the possible values of $l = 0, 1, 2$. The possible values of m will be $\pm l$ that are $-2, -1, 0, 1, 2$ and s will always be $\pm \frac{1}{2}$.

The value of m will never more than l , thus this is not the valid set of quantum numbers.

32. If x is the speed of electron in the Bohr's first orbit of hydrogen atom, then find the speed of electron in the fourth Bohr's orbit

(a) $8x$ (b) $x/4$

(c) $4x$ (d) $x/8$

Answer: (c)

According to Bohr's model the velocity of an electron for hydrogen and hydrogen like atoms is quantised and is equals $(\frac{2.18 \times 10^6 \times Z}{n})$.

Here, n and Z are orbit number and atomic number respectively. For H atom Z = 1. The velocity of electron in hydrogen atom's 1 orbit = $2.18 \times 10^6 = x$

The velocity of electron in hydrogen atom's 4 orbit = $2.18 \times 10^6 \times 4 = 4x$

Hence, the velocity in fourth bohr orbit is 4x.

33. Choose the correct set of quantum number for the Ne atom's electron of second excited state

(a) 2,0,0,±1/2

(b) 3,2,1,±1/2

(c) 3,0,0,±1/2

(d) 3,1,2,±1/2

Answer: (c)

Ground state electronic configuration of Ne atom is $1s^2, 2s^2, 2p^6$.

Electronic Configuration for its first excitation state $1s^2, 2s^2, 2p^5, 3s^1$.

Electronic Configuration for its second excitation state $1s^2, 2s^2, 2p^4, 3s^2$.

Quantum numbers for 3 electrons will be 3,0,0,± $\frac{1}{2}$.

34. Select the one among the following transitions in hydrogen atom, which gives an absorption line of lowest frequency

(a) $n = 1$ to $n = 3$

(b) $n = 2$ to $n = 3$

(c) $n = 4$ to $n = 6$

(d) $n = 3$ to $n = 4$

Answer: (c)

When energy is absorbed, electron shifts from lower to higher orbit and the absorption line in the spectra arises. $n = 4$ to $n = 6$ will have the lowest frequency as this falls in the bracket series.

35. An electron is moving with a kinetic energy of $4.55 \times 10^{-25} \text{ J}$.

What will be the de Broglie wave length for this electron?

(a) $5.28 \times 10^{-7} \text{ m}$

(b) $7.28 \times 10^{-7} \text{ m}$

(c) $2 \times 10^{-10} \text{ m}$

(d) $3 \times 10^{-5} \text{ m}$

Answer: (b)

$$\text{KE} = \frac{1}{2}mv^2 = 4.55 \times 10^{-25}$$

$$v^2 = \frac{2 \times 4.55 \times 10^{-25}}{9.1 \times 10^{-31}} = 1 \times 10^6$$

$$v = 103 \text{ m/s}$$

Answer: (b)

From Einstein's equation for photoelectric effect

$$h\nu = h\nu_0 + \frac{1}{2}mv^2$$

$$\Rightarrow \frac{1}{2}mv^2 = h\nu - h\nu_0$$

$$\Rightarrow \frac{1}{2}mv^2 = 2h\nu_0 - h\nu_0 = h\nu_0 \quad \text{-----(1)}$$

$$\frac{1}{2}mv^2 = 5h\nu_0 - h\nu_0 = 4h\nu_0 \quad \text{-----(2)}$$

$$\frac{\text{Eqn(1)}}{\text{Eqn(2)}} \Rightarrow \frac{v^2}{v^2} = \frac{1}{4} \Rightarrow \frac{v_1}{v_2} = \frac{1}{2}$$

38. Number of waves in the third Bohr's orbit of hydrogen will be

(a) 3

(b) 6

(c) 9

(d) 12

Answer: (a)

Number of waves = $\frac{\text{circumference}}{\text{wavelength}} = \frac{2\pi r}{\lambda}$

$$= \frac{2\pi r}{h/mv} = \frac{2\pi}{h} (mvr)$$

$$= \frac{2\pi}{h} \times \frac{nh}{2\pi}$$

$$n = 3$$

Therefore, the electronic transition involved should be $n_2 = \infty \rightarrow n_1 = 2$ (as it belongs to the Balmer series)

$$\frac{1}{\lambda} = RZ^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$$\frac{1}{\lambda_{\text{shortest}}} = 109678 \times 12^2 \left(\frac{1}{2^2} - \frac{1}{\infty^2} \right)$$

$$\lambda = 3.647 \times 10^{-5} \text{ cm}$$

$$= 3647 \text{ \AA}$$

41. For a d-electron, the orbital angular momentum is:

- (a) $6\hbar$ (B) $2\hbar$ (C) \hbar (D) $2\hbar$

Answer: (a)

For d-electron, $l=2$, orbital angular momentum

$$= \sqrt{l(l+1)}\hbar = \sqrt{2(2+1)}\hbar = \sqrt{6}\hbar$$

Hence, (a) is the correct answer.

42. The electronic configuration of an element is $1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 3d^5, 4s^1$. This represents its

- (A) excited state (B) ground state

(C) cationic form

(D) anionic form

Answer: (b)

The given electronic configuration is ground state for chromium.

Hence, (B) is the correct answer.

(A) $\text{Cr}^{3+}, \text{Fe}^{3+}$ (B) $\text{Fe}^{3+}, \text{Mn}^{2+}$ (C) $\text{Fe}^{3+}, \text{Co}^{3+}$ (D) $\text{Se}^{3+}, \text{Cr}^{3+}$

Answer: (b)

Fe^{3+} and Mn^{2+} have same electronic configuration

Hence (B) is the correct answer.

44. Principal, azimuthal and magnetic quantum numbers are respectively related to

(A) size, orientation and shape

(B) size shape and orientation

(C) shape, size and orientation

(D) none of these

Answer: (b)

Principal gives, i.e. azimuthal gives shape and magnetic quantum number gives the orientation

Hence, (B) is the correct answer.

45. Number of waves in third Bohr's orbit of hydrogen will be

- (A) 3 (B) 6 (C) 9 (D) 12

Answer: (a)

Number of waves = $\frac{\text{Circumference}}{\text{Wavelength}}$

$$\frac{2\pi r}{\lambda} = \frac{2\pi r}{h/mv} = \frac{2\pi}{h}(mvr)$$

$$= \frac{2\pi}{h} \times \frac{nh}{2\pi}$$

$$\therefore n = 3$$

Hence, (A) is the correct answer.