

**FINAL JEE(Advanced) EXAMINATION - 2022**  
(Held On Sunday 28th AUGUST, 2022)

PAPER-1

TEST PAPER WITH SOLUTION

## CHEMISTRY

### SECTION-1 : (Maximum Marks : 24)

- This section contains **EIGHT (08)** questions. The answer to each question is a **NUMERICAL VALUE**. For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places. Answer to each question will be evaluated according to the following marking scheme.

*Full Marks* : +3 **ONLY** if the correct numerical value is entered;  
*Zero Marks* : 0 In all other cases.

1. 2 mol of Hg(g) is combusted in a fixed volume bomb calorimeter with excess of O<sub>2</sub> at 298 K and 1 atm into HgO(s). During the reaction, temperature increases from 298.0 K to 312.8 K. If heat capacity of the bomb calorimeter and enthalpy of formation of Hg(g) are 20.00 kJ K<sup>-1</sup> and 61.32 kJ mol<sup>-1</sup> at 298 K, respectively, the calculated standard molar enthalpy of formation of HgO(s) at 298 K is X kJ mol<sup>-1</sup>. The value of |X| is \_\_\_\_\_.  
[Given : Gas constant R = 8.3 J K<sup>-1</sup> mol<sup>-1</sup>]


**Ans. (90.39)**

**Sol.** Q<sub>rxn</sub> = CΔT

$$|Q| \times 2 = 20 \times 14.8$$


$$|Q| = 148 \text{ kJ/mol}$$

$$Q = -148 \text{ kJ/mol}$$



$$Q_H = Q_U + \Delta n_g RT$$

$$= -148 - \frac{3}{2} \times \frac{8.3}{1000} \times 298 = -151.7101$$



$$Q_H = -151.7101 + 61.32 = -90.39 \text{ kJ/mol}$$

Ans. 90.39




5. Dissolving 1.24 g of white phosphorous in boiling NaOH solution in an inert atmosphere gives a gas **Q**. The amount of CuSO<sub>4</sub> (in g) required to completely consume the gas **Q** is \_\_\_\_\_.  
[Given : Atomic mass of H = 1, O = 16, Na = 23, P = 31, S = 32, Cu = 63]


Ans. (2.38 / 2.39)

Sol. Mole of P<sub>4</sub> =

$$\frac{1.24}{31 \times 4} = 0.01$$



$$0.01 \text{ mole} \qquad \qquad 0.01 \text{ mole}$$




$$0.01 \quad \frac{3}{2} \times 0.01$$

$$= \frac{0.03}{2} \text{ moles}$$

$$W_{CuSO_4} = \frac{0.03}{2} \times 159 = 2.385 \text{ gm}$$

Ans. = 2.38 or 2.39


6. Consider the following reaction.



On estimation of bromine in 1.00 g of **R** using Carius method, the amount of AgBr formed (in g) is

[Given : Atomic mass of H = 1, C = 12, O = 16, P = 31, Br = 80, Ag = 108]

Ans. (1.50)




$$1gR \rightarrow \frac{1}{250} \text{ moles}$$

$$\text{No. of Br Atoms} \rightarrow \frac{2}{250} \text{ moles}$$

$$\text{Moles of AgBr} \rightarrow \frac{2}{250} \text{ moles}$$


$$\text{Mass of AgBr} = \frac{2}{250} \times (108 + 80) = 1.504$$

7. The weight percentage of hydrogen in **Q**, formed in the following reaction sequence, is \_\_\_\_\_.




[Given : Atomic mass of H = 1, C = 12, N = 14, O = 16, S = 32, Cl = 35]

Ans. (1.31)



8. If the reaction sequence given below is carried out with 15 moles of acetylene, the amount of the product **D** formed (in g) is \_\_\_\_\_.




The yields of **A**, **B**, **C** and **D** are given in parentheses.

[Given : Atomic mass of H = 1, C = 12, O = 16, Cl = 35]

Ans. (136)

Sol.



**SECTION-2 : (Maximum Marks : 24)**

□□ This section contains **SIX (06)** questions.  
 □□ Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).

□□ For each question, choose the option(s) corresponding to (all) the correct answer(s).  
 □ Answer to each question will be evaluated according to the following marking scheme:

Full Marks	: +4 <b>ONLY</b> if (all) the correct option(s) is(are) chosen;
Partial Marks	: +3 If all the four options are correct but <b>ONLY</b> three options are chosen;
Partial Marks	: +2 If three or more options are correct but <b>ONLY</b> two options are chosen, both of which are correct;
Partial Marks	: +1 If two or more options are correct but <b>ONLY</b> one option is chosen and it is a correct option;
Zero Marks	: 0 If none of the options is chosen (i.e. the question is unanswered);
Negative Marks	: -2 In all other cases.

9. For diatomic molecules, the correct statement(s) about the molecular orbitals formed by the overlap to two  $2p_z$  orbitals is(are)


(A)  $\sigma$  orbital has a total of two nodal planes.


(B)  $\sigma^*$  orbital has one node in the  $xz$ -plane containing the molecular axis.

(C)  $\pi$  orbital has one node in the plane which is perpendicular to the molecular axis and goes through the center of the molecule.

(D)  $\pi^*$  orbital has one node in the  $xy$ -plane containing the molecular axis.

Ans. (A,D)





10. The correct option(s) related to adsorption processes is(are)

- (A) Chemisorption results in a unimolecular layer.
- (B) The enthalpy change during physisorption is in the range of 100 to 140 kJ mol<sup>-1</sup>
- (C) Chemisorption is an endothermic process.
- (D) Lowering the temperature favors physisorption processes.

Ans. (A,D)

Sol. (A) Chemisorption is unimolecular layered.

- (B) Enthalpy of physisorption is much less in magnitude.
- (C) Chemisorption of gases on solids is exothermic.
- (D) As physisorption is exothermic so lowering temperature favours it.

11. The electrochemical extraction of aluminum from bauxite ore involves.


- (A) the reaction of Al<sub>2</sub>O<sub>3</sub> with coke (C) at a temperature > 2500°C.
- (B) the neutralization of aluminate solution by passing CO<sub>2</sub> gas to precipitate hydrated alumina (Al<sub>2</sub>O<sub>3</sub>·3H<sub>2</sub>O)
- (C) the dissolution of Al<sub>2</sub>O<sub>3</sub> in hot aqueous NaOH.
- (D) the electrolysis of Al<sub>2</sub>O<sub>3</sub> mixed with Na<sub>3</sub>AlF<sub>6</sub> to give Al and CO<sub>2</sub>.

Ans. (B,C,D)

Sol. (A) Electrochemical extraction of Aluminum from bauxite done below 2500°C



The sodium aluminate present in solution is neutralised by passing CO<sub>2</sub> gas and hydrated Al<sub>2</sub>O<sub>3</sub> is precipitated.



Concentration of bauxite is carried out by heating the powdered ore with hot concentrated solution of NaOH

(D) In metallurgy of aluminum, Al<sub>2</sub>O<sub>3</sub> is mixed with Na<sub>3</sub>AlF<sub>6</sub>


12. The treatment of galena with  $\text{HNO}_3$  produces a gas that is
- (A) paramagnetic (B) bent in geometry  
(C) an acidic oxide (D) colorless

Ans. (A,D)

Sol.  $3\text{PbS} + 8\text{HNO}_3 \rightarrow 3\text{Pb}(\text{NO}_3)_2 + 2\text{NO} + 4\text{H}_2\text{O} + \text{S}$


$\text{NO}$  □ Neutral oxide, Paramagnetic, Linear geometry, Colourless gas

13. Considering the reaction sequence given below, the correct statement(s) is(are)




- (A) **P** can be reduced to a primary alcohol using  $\text{NaBH}_4$ .  
 (B) Treating **P** with conc.  $\text{NH}_4\text{OH}$  solution followed by acidification gives **Q**.  
 (C) Treating **Q** with a solution of  $\text{NaNO}_2$  in aq.  $\text{HCl}$  liberates  $\text{N}_2$ .  
 (D) **P** is more acidic than  $\text{CH}_3\text{CH}_2\text{COOH}$ .

Ans. (B,C,D)



Sol.

14. Consider the following reaction sequence,



the correct option(s) is(are)

(A) P = H<sub>2</sub>/Pd, ethanol




R = NaNO<sub>2</sub>/HCl

U = 1. H<sub>3</sub>PO<sub>2</sub>

2. KMnO<sub>4</sub> - KOH, heat


(B) P = Sn/HCl

R = HNO<sub>2</sub>




U = 1. CH<sub>3</sub>CH<sub>2</sub>OH

2. KMnO<sub>4</sub> - KOH, heat




R = H<sub>2</sub>/Pd, ethanol



Ans. (A,B,C)

Sol.










**SECTION-3 : (Maximum Marks : 12)**

- This section contains **FOUR (04)** Matching List Sets.  
 Each set has **ONE** Multiple Choice Question.  
 Each set has **TWO** lists : **List-I** and **List-II**.  
 **List-I** has **Four** entries (I), (II), (III) and (IV) and **List-II** has **Five** entries (P), (Q), (R), (S) and (T).  
 **FOUR** options are given in each Multiple Choice Question based on **List-I** and **List-II** and **ONLY ONE** of these four options satisfies the condition asked in the Multiple Choice Question.  
 Answer to each question will be evaluated according to the following marking scheme:  
*Full Marks* : +3 **ONLY** if the option corresponding to the correct combination is chosen;  
*Zero Marks* : 0 If none of the options is chosen (i.e. the question is unanswered);  
*Negative Marks* : -1 In all other cases.

**15.** Match the rate expressions in LIST-I for the decomposition of X with the corresponding profiles provided in LIST-II. Xs and k constants having appropriate units.

LIST-I	LIST-II
(I) $\text{rate} = \frac{k[X]^2}{X_s + X}$ under all possible initial concentration of X	(P) 
(II) $\text{rate} = \frac{k[X]}{X_s + X}$ where initial concentration of X are much less than Xs	(Q) 
(III) $\text{rate} = \frac{k[X]^2}{X_s + X}$ where initial concentration of X are much higher than Xs	(R) 

<p>(IV)</p> $\text{rate} = \frac{k[X]^2}{X_s - [X]}$ <p>where initial concentration of X is much higher than <math>X_s</math></p>	<p>(S)</p> 
	<p>(T)</p> 

- (A) I □ P; II □ Q; III □ S; IV □ T  
 (B) I □ R; II □ S; III □ S; IV □ T  
 (C) I □ P; II □ Q; III □ Q; IV □ R  
 (D) I □ R; II □ S; III □ Q; IV □ R

**Ans. (A)**

**Sol. (I)** 
$$\text{rate} = \frac{k[x]}{x_s - [x]} = \frac{k}{\frac{x_s}{[x]} - 1}$$

If  $[x] \gg x_s$  rate =  $k$  □ order = 0

□ (I) – (R), (P)

(II)  $[x] < x_s$  □ rate =  $\frac{k[x]}{x_s}$  □ order = 1

□ (II) – (Q), (T)

(III)  $[x] \gg x_s$  □ rate =  $k$  □ order = 0

□ (III) – (P), (S)

(IV) rate =  $\frac{k[x]^2}{x_s - [x]}$

$[x] \gg x_s$  □ rate =  $k[x]$

□ (IV) – (Q), (T)

**Ans. (A)**

16. LIST-I contains compounds and LIST-II contains reaction

LIST-I

- (I)  $\text{H}_2\text{O}_2$   
 (II)  $\text{Mg}(\text{OH})_2$   
 (III)  $\text{BaCl}_2$   
 (IV)  $\text{CaCO}_3$

LIST-II

- (P)  $\text{Mg}(\text{HCO}_3)_2 + \text{Ca}(\text{OH})_2 \rightarrow$   
 (Q)  $\text{BaO}_2 + \text{H}_2\text{SO}_4 \rightarrow$   
 (R)  $\text{Ca}(\text{OH})_2 + \text{MgCl}_2 \rightarrow$   
 (S)  $\text{BaO}_2 + \text{HCl} \rightarrow$   
 (T)  $\text{Ca}(\text{HCO}_3)_2 + \text{Ca}(\text{OH})_2 \rightarrow$

Match each compound in LIST – I with its formation reaction(s) in LIST-II, and choose the correct option

- (A) I  $\rightarrow$  Q; II  $\rightarrow$  P; III  $\rightarrow$  S; IV  $\rightarrow$  R  
 (B) I  $\rightarrow$  T; II  $\rightarrow$  P; III  $\rightarrow$  Q; IV  $\rightarrow$  R  
 (C) I  $\rightarrow$  T; II  $\rightarrow$  R; III  $\rightarrow$  Q; IV  $\rightarrow$  P  
 (D) I  $\rightarrow$  Q; II  $\rightarrow$  R; III  $\rightarrow$  S; IV  $\rightarrow$  P

**Ans. (D)**

**Sol.** (P)  $\text{Mg}(\text{HCO}_3)_2 + 2\text{Ca}(\text{OH})_2 \rightarrow \text{Mg}(\text{OH})_2 + 2\text{CaCO}_3 + 2\text{H}_2\text{O}$

(Q)  $\text{BaO}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{H}_2\text{O}_2 + \text{BaSO}_4$

(R)  $\text{Ca}(\text{OH})_2 + \text{MgCl}_2 \rightarrow \text{Mg}(\text{OH})_2 + \text{CaCl}_2$

(S)  $\text{BaO}_2 + 2\text{HCl} \rightarrow \text{BaCl}_2 + \text{H}_2\text{O}_2$

(T)  $\text{Ca}(\text{HCO}_3)_2 + \text{Ca}(\text{OH})_2 \rightarrow 2\text{CaCO}_3 + 2\text{H}_2\text{O}$

17. LIST-I contains metal species and LIST-II contains their properties.

LIST-I

- (I)  $[\text{Cr}(\text{CN})_6]^{4-}$   
 (II)  $[\text{RuCl}_6]^{2-}$   
 (III)  $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$   
 (IV)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$

LIST-II

- (P) f2g orbitals contain 4 electrons  
 (Q)  $\mu_{\text{spin-only}} = 4.9 \text{ BM}$   
 (R) low spin complex ion  
 (S) metal ion in  $4+$  oxidation state

(T)  $d^4$  species

[Given : Atomic number of Cr = 24, Ru = 44, Fe = 26]


Match each metal species in LIST-I with their properties in LIST-II, and choose the correct option

- (A) I  $\rightarrow$  R, T; II  $\rightarrow$  P, S; III  $\rightarrow$  Q, T; IV  $\rightarrow$  P, Q  
 (B) I  $\rightarrow$  R, S; II  $\rightarrow$  P, T; III  $\rightarrow$  P, Q; IV  $\rightarrow$  Q, T  
 (C) I  $\rightarrow$  P, R; II  $\rightarrow$  R, S; III  $\rightarrow$  R, T; IV  $\rightarrow$  P, T  
 (D) I  $\rightarrow$  Q, T; II  $\rightarrow$  S, T; III  $\rightarrow$  P, T; IV  $\rightarrow$  Q, R

**Ans. (A)**

Sol. (1)  $[\text{Cr}(\text{CN})_6]^{4-}$


$\text{Cr}^{+2} [\text{Ar}]_{18} 3d^4 4s^0$ , low spin complex



P,R,T

(2)  $[\text{RuCl}_6]^{2-}$


$\text{Ru}^{+4} [\text{Kr}]_{36} 4d^4 5s^0$ , low spin complex



P,R,S,T

(3)  $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$


$\text{Cr}^{+2} [\text{Ar}]_{18} 3d^4 4s^0$ , high spin complex



Q,T

(4)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$

$\text{Fe}^{+2} [\text{Ar}]_{18} 3d^6 4s^0$ , high spin complex



P,Q

18. Match the compounds in LIST-I with the observation in LIST-II, and choose the correct option.

LIST-I

(I) Aniline

(II) o-Cresol

(III) Cysteine

LIST-II

(P) Sodium fusion extract of the compound on boiling with  $\text{FeSO}_4$ , followed by acidification with conc.  $\text{H}_2\text{SO}_4$ , gives Prussian blue color.


(Q) Sodium fusion extract of the compound on treatment with sodium nitroprusside gives blood red color.


(R) Addition of the compound to a saturated solution of  $\text{NaHCO}_3$  results in effervescence.

(IV) Coprolactam

(S) The compound reacts with bromine water to give a white precipitate.

(T) Treating the compound with neutral  $\text{FeCl}_3$  solution produces violet color.(A) I  P, Q; II  S; III  Q, R; IV  P(B) I  P; II  R, S; III  R; IV  Q, S(C) I  Q, S; II  P, T; III  P; IV  S(D) I  P, S; II  T; III  Q, R; IV  P**Ans. (D)**

**Sol.**  : Blue colour in Lassaign test due to presence of N  
Aniline

 : Violet colour with  $\text{FeCl}_3$  due to presence of phenolic OH  
o-Cresol

$\text{HS}-\text{CH}_2-\underset{\text{NH}_2}{\text{CH}}-\text{COOH}$   
Cystein : It gives blood red colour with  $\text{NaSCN}$

 : Blue colour in Lassaign test due to presence of N  
Caprolactam